# PRUXIENCE® Patient Enrolment, Rx and Consent Form for Rheumatoid Arthritis

Please fax to 1-833-958-3539 or 1-833-958-FLEX upon completion. To speak to a clinic representative, call toll-free 1-855-935-3539 or 1-855-935-FLEX.

PATIENT INFORMATION		<b>PHYSICIAN</b>	INFORMATION	
Name:		Physician Name:		
Address:		Address:		
<b>Tel.</b> (Home): <b>Tel.</b> (Other):				
Okay to leave messages:  Yes No		Tel (Office):		
Best time to be contacted:   AM PM Night		Fax (Office):		
Email: Date of birth: DD/MM/YYYY		, ,		
Known allergies: No Yes If Yes, please specify:				
Triown allergies. — NO — res il res, please specify:		Cilidii.		
PHYSICIAN PRESCRIBING SECTION Please consult the RUXIE	NCE® Broduct Manager	anh for important info	ormation relating to decing administration adverse reactions and drug in	toractions
				teractions.
Confirmed diagnosis of: Moderately to severely active Rheumatoid Arthritis (RA)				
ORDER FOR RUXIENCE®				
☐ First treatment ☐ Subsequent treatment			TREATMENT LEGEND	
Anticipated infusion date: DD/MM/YYYY Infusion clinic location:  First treatment:			<b>255 minute (4.25 hrs):</b> RUXIENCE® 1000 mg IV at a rate of 50 mg/hr for 30 minutes, increasing 50 mg/hr every 30 minutes as tolerated, for a maxil 400 mg/hr.	
<b>Day 1</b> ☐ 255 minute infusion (4.25 hrs) × 1000 mg			195 minute (3.25 hrs): RUXIENCE® 1000 mg IV can be started at a rate of	of 100 mg/hr
Day 15 ☐ 195 minute infusion (3.25 hrs) × 1000 mg OR			for the first 30 minutes, increasing 100 mg/hr every 30 minutes as tolerated, for a	
☐ Alternative 120 minute infusion (2 hrs) × 1000 mg*			maximum rate of 400 mg/hr.	
Subsequent treatments:			120 minute (2 hrs): If patients did not experience a serious infusion-rela	
Day 1 ☐ 255 minute infusion (4.25 hrs) × 1000 mg OR			event during the previous infusion administered using the standard admi schedule, an alternative 120-minute infusion of a concentration at 4 mg/r	
Alternative 120 minute infusion (2 hrs) × 1000 mg	*		volume can be administered for the second infusion. Initiate at a rate of 6	2.5 mL/hour
<b>Day 15</b> 195 minute infusion (3.25 hrs) × 1000 mg <b>OR</b>			(125 mg) given in the first 30 minutes and 150 mL/hour (875 mg) given	
Alternative 120 minute infusion (2 hrs) × 1000 mg			90 minutes. If the 120-minute infusion is tolerated, the same alternative infusion rate can be used when administering subsequent infusions and	
* Alternative 120 minute infusion is not an option for all patie	ents. Consult the	Product	an option for all patients. Consult the Product Monograph for information	
Monograph for information on alternative administration e	ligibility.		administration eligibility.	
U Other dosing:				
-				
☐ Dilute RUXIENCE® to a final concentration of 1 to 4 mg/mL	Other instruction	ons:	Comments:	
into an infusion bag containing either 0.9% Sodium				
<u>-</u>			Comments:	
into an infusion bag containing either 0.9% Sodium	Blood pressure	meds on hold		
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP	Blood pressure	meds on hold	I: YES NO Please specify:INFUSION REACTIONS	
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion	Blood pressure PRN MEDICA  Acetaminop	meds on hold ATIONS FOR when 325-650 n	I: NO Please specify:	
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion Diphenhydramine 50 mg PO 15-30 min pre infusion	Blood pressure PRN MEDICA Acetaminop Dimenhydri	e meds on hold ATIONS FOR Then 325-650 n nate 25-50 mg	I: YES NO Please specify:	
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL	Blood pressure PRN MEDICA Acetaminop Dimenhydri Diphenhydri Epinephrine	e meds on hold ATIONS FOR Then 325-650 m Then 325-50 mg Tamine 25-50 mg	I: YES NO Please specify:	itus, hives
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min	Blood pressure PRN MEDICA Acetaminop Dimenhydri Diphenhydri Epinephrine severe anap	e meds on hold ATIONS FOR when 325-650 m nate 25-50 mg ramine 25-50 m e (1:1000) 0.01 ohylactic reactio	I: YES NO Please specify:	itus, hives
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min prior to infusion	Blood pressure PRN MEDICA Acetaminop Dimenhydri Diphenhydr Epinephrine severe anap Hydrocortise	e meds on hold ATIONS FOR when 325-650 m nate 25-50 mg ramine 25-50 m e (1:1000) 0.01 whylactic reactione 100 mg IV	I: YES NO Please specify:  INFUSION REACTIONS  INFUSION REACTIONS  INFUSION Q 4-6 hours, for pain and fever, chills INFO/IV PRN Q 4-6 hours, for nausea and vomiting INFO/IV PRN Q 4-6 hours for itching, urticaria, pruse INFO/IV PRN Q 10-15 minutes × 20 INFO/IV PRN × 1 for severe allergic/anaphylactic reaction	itus, hives
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min prior to infusion  Other (indicate):	Blood pressure  PRN MEDICA  Acetaminop  Dimenhydri  Diphenhydr  Epinephrine severe anap  Hydrocortise  Oxygen via	e meds on hold ATIONS FOR when 325-650 m nate 25-50 mg amine 25-50 mg (1:1000) 0.01 shylactic reactione 100 mg IV mask/nasal pro	I: YES NO Please specify:  INFUSION REACTIONS  INFUSION Q 4-6 hours, for pain and fever, chills INFO/IV PRN Q 4-6 hours, for nausea and vomiting INFO/IV PRN Q 4-6 hours for itching, urticaria, pruse INFO/IV PRN Q 10-15 minutes × 2 INFO/IV PRN × 1 for severe allergic/anaphylactic reaction INFO/IV PRN × 1 for shortness of breath, wheezing	itus, hives
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min prior to infusion  Other (indicate):  Pre-medications not required	Blood pressure PRN MEDICA Acetaminop Dimenhydri Diphenhydri Epinephrine severe anap Hydrocortise Oxygen via Salbutamol	e meds on hold ATIONS FOR Then 325-650 mg Then 325-50 mg Tamine 25-50 mg Tamin	I: YES NO Please specify:  INFUSION REACTIONS  INFUSION Q 4-6 hours, for pain and fever, chills INFO/IV PRN Q 4 hours, for nausea and vomiting INFO/IV/IM PRN Q 4-6 hours for itching, urticaria, prunication INFO/IM PRN Q 10-15 minutes × 2000 INFO/IM PRN Y 10-15 minutes × 2000 INFO/IM PRN Y 1 for severe allergic/anaphylactic reaction INFO/IM PRN FOR STANDARD FOR STANDA	itus, hives
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min prior to infusion  Other (indicate):	Blood pressure PRN MEDICA Acetaminop Dimenhydri Diphenhydri Epinephrine severe anap Hydrocortise Oxygen via Salbutamol Other (indice	e meds on hold ATIONS FOR Then 325-650 mg Then 325-50 mg Tamine 25-50 mg Tamin	I: YES NO Please specify:  INFUSION REACTIONS  Ing PO PRN q 4-6 hours, for pain and fever, chills and PO/IV PRN q 4 hours, for nausea and vomiting ang PO/IV/IM PRN q 4-6 hours for itching, urticaria, pruntly (max. 0.5 mL) SC/IM PRN q 10-15 minutes × 2 con PRN × 1 for severe allergic/anaphylactic reaction angs PRN for shortness of breath, wheezing ours via aerochamber PRN for dyspnea, wheezing	itus, hives
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min prior to infusion  Other (indicate):  Pre-medications not required  Please specify:	Blood pressure PRN MEDICA Acetaminop Dimenhydri Diphenhydri Epinephrine severe anap Hydrocortise Oxygen via Salbutamol Other (indice	e meds on hold ATIONS FOR Then 325-650 mg Then 325-50 mg Tamine 25-50 mg Tamin	I: YES NO Please specify:  INFUSION REACTIONS  INFUSION Q 4-6 hours, for pain and fever, chills INFO/IV PRN Q 4 hours, for nausea and vomiting INFO/IV/IM PRN Q 4-6 hours for itching, urticaria, prunication INFO/IM PRN Q 10-15 minutes × 2000 INFO/IM PRN Y 10-15 minutes × 2000 INFO/IM PRN Y 1 for severe allergic/anaphylactic reaction INFO/IM PRN FOR STANDARD FOR STANDA	itus, hives
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min prior to infusion  Other (indicate):  Pre-medications not required  Please specify:  PHYSICIAN SIGNATURE	Blood pressure  PRN MEDICA  Acetaminop  Dimenhydri  Diphenhydri  Epinephrine severe anap  Hydrocortise  Oxygen via  Salbutamol  Other (indic	e meds on hold ATIONS FOR when 325-650 m nate 25-50 mg (1:1000) 0.01 hylactic reaction one 100 mg IV mask/nasal pro 2 puffs q 4-6 ho nate):	I: YES NO Please specify:  INFUSION REACTIONS  INFUSION Q 4-6 hours, for pain and fever, chills INFO/IV PRN Q 4-6 hours, for nausea and vomiting INFO/IV PRN Q 4-6 hours for itching, urticaria, prunchly (max. 0.5 mL) SC/IM PRN Q 10-15 minutes × 2000  PRN × 1 for severe allergic/anaphylactic reaction longs PRN for shortness of breath, wheezing ours via aerochamber PRN for dyspnea, wheezing	itus, hives for
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min prior to infusion  Other (indicate):  Pre-medications not required  Please specify:	Blood pressure  PRN MEDICA  Acetaminop  Dimenhydri  Diphenhydri  Epinephrine severe anap  Hydrocortise  Oxygen via  Salbutamol  Other (indic	e meds on hold ATIONS FOR when 325-650 m nate 25-50 mg (1:1000) 0.01 hylactic reaction one 100 mg IV mask/nasal pro 2 puffs q 4-6 ho nate):	I: YES NO Please specify:  INFUSION REACTIONS  INFUSION Q 4-6 hours, for pain and fever, chills INFO/IV PRN Q 4-6 hours, for nausea and vomiting INFO/IV PRN Q 4-6 hours for itching, urticaria, prunchly (max. 0.5 mL) SC/IM PRN Q 10-15 minutes × 2000  PRN × 1 for severe allergic/anaphylactic reaction longs PRN for shortness of breath, wheezing ours via aerochamber PRN for dyspnea, wheezing	itus, hives for
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min prior to infusion  Other (indicate):  Pre-medications not required  Please specify:  PHYSICIAN SIGNATURE	Blood pressure  PRN MEDICA  Acetaminop  Dimenhydri  Diphenhydri  Epinephrine severe anap  Hydrocortise  Oxygen via  Salbutamol  Other (indice  PRN medica	e meds on hold ATIONS FOR when 325-650 m nate 25-50 mg amine 25-50 mg (1:1000) 0.01 ohylactic reaction one 100 mg IV mask/nasal pro 2 puffs q 4-6 hold rate):	I: YES NO Please specify:  INFUSION REACTIONS  INFUSION Q 4-6 hours, for pain and fever, chills INFO/IV PRN Q 4-6 hours, for nausea and vomiting INFO/IV PRN Q 4-6 hours for itching, urticaria, prunchly (max. 0.5 mL) SC/IM PRN Q 10-15 minutes × 2000  PRN × 1 for severe allergic/anaphylactic reaction longs PRN for shortness of breath, wheezing ours via aerochamber PRN for dyspnea, wheezing	itus, hives for
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min prior to infusion  Other (indicate):  Pre-medications not required  Please specify:  PHYSICIAN SIGNATURE  Do you accept that Pfizer Canada's Drug Safety Unit contact you results to the property of the p	Blood pressure PRN MEDICA Acetaminop Dimenhydri Diphenhydri Epinephrine severe anap Hydrocortise Oxygen via Salbutamol Other (indice PRN medica	e meds on hold ATIONS FOR when 325-650 mg amine 25-50 mg amine 25-50 mg (1:1000) 0.01 whylactic reaction one 100 mg IV mask/nasal pro 2 puffs q 4-6 ho atel):	I: YES NO Please specify:  INFUSION REACTIONS  Ing PO PRN q 4-6 hours, for pain and fever, chills IPO/IV PRN q 4 hours, for nausea and vomiting INFUSION Q 4-6 hours for itching, urticaria, prunchly (max. 0.5 mL) SC/IM PRN q 10-15 minutes × 2 con INFUSION PRN × 1 for severe allergic/anaphylactic reaction longs PRN for shortness of breath, wheezing ours via aerochamber PRN for dyspnea, wheezing  INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever hours fo	) NO le reverse.
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min prior to infusion  Other (indicate):  Pre-medications not required  Please specify:  PHYSICIAN SIGNATURE  Do you accept that Pfizer Canada's Drug Safety Unit contact you results.	Blood pressure PRN MEDICA Acetaminop Dimenhydri Diphenhydri Epinephrine severe anap Hydrocortise Oxygen via Salbutamol Other (indice PRN medica	e meds on hold ATIONS FOR when 325-650 mg amine 25-50 mg amine 25-50 mg (1:1000) 0.01 whylactic reaction one 100 mg IV mask/nasal pro 2 puffs q 4-6 ho atel):	I: YES NO Please specify:  INFUSION REACTIONS  Ing PO PRN q 4-6 hours, for pain and fever, chills IPO/IV PRN q 4 hours, for nausea and vomiting INFUSION Q 4-6 hours for itching, urticaria, prunchly (max. 0.5 mL) SC/IM PRN q 10-15 minutes × 2 con INFUSION PRN × 1 for severe allergic/anaphylactic reaction longs PRN for shortness of breath, wheezing ours via aerochamber PRN for dyspnea, wheezing  INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever hours fo	) NO le reverse.
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min prior to infusion  Other (indicate):  Pre-medications not required  Please specify:  PHYSICIAN SIGNATURE  Do you accept that Pfizer Canada's Drug Safety Unit contact you results to the property of the p	Blood pressure PRN MEDICA Acetaminop Dimenhydri Diphenhydri Epinephrine severe anap Hydrocortise Oxygen via Salbutamol Other (indice PRN medica	e meds on hold ATIONS FOR when 325-650 mg amine 25-50 mg amine 25-50 mg (1:1000) 0.01 whylactic reaction one 100 mg IV mask/nasal pro 2 puffs q 4-6 ho atel):	I: YES NO Please specify:  INFUSION REACTIONS  Ing PO PRN q 4-6 hours, for pain and fever, chills IPO/IV PRN q 4 hours, for nausea and vomiting INFUSION Q 4-6 hours for itching, urticaria, prunchly (max. 0.5 mL) SC/IM PRN q 10-15 minutes × 2 con INFUSION PRN × 1 for severe allergic/anaphylactic reaction longs PRN for shortness of breath, wheezing ours via aerochamber PRN for dyspnea, wheezing  INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever hours fo	) NO le reverse.
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min prior to infusion  Other (indicate):  Pre-medications not required  Please specify:  PHYSICIAN SIGNATURE  Do you accept that Pfizer Canada's Drug Safety Unit contact your Notes:  SIGN HERE:  † Effective date. Order(s) expires one year from the date of signature. Prescriber certificat	Blood pressure PRN MEDICA Acetaminop Dimenhydri Diphenhydri Epinephrine severe anap Hydrocortise Oxygen via Salbutamol Other (indice PRN medica	e meds on hold ATIONS FOR when 325-650 mg amine 25-50 mg amine 25-50 mg (1:1000) 0.01 whylactic reaction one 100 mg IV mask/nasal pro 2 puffs q 4-6 ho atel):	I: YES NO Please specify:  INFUSION REACTIONS  Ing PO PRN q 4-6 hours, for pain and fever, chills IPO/IV PRN q 4 hours, for nausea and vomiting INFUSION Q 4-6 hours for itching, urticaria, prunchly (max. 0.5 mL) SC/IM PRN q 10-15 minutes × 2 con INFUSION PRN × 1 for severe allergic/anaphylactic reaction longs PRN for shortness of breath, wheezing ours via aerochamber PRN for dyspnea, wheezing  INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours, for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever, chills INFUSION PRN q 4-6 hours for pain and fever hours fo	NO e reverse.
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min prior to infusion  Other (indicate):  Pre-medications not required  Please specify:  PHYSICIAN SIGNATURE  Do you accept that Pfizer Canada's Drug Safety Unit contact you result to the contact your street of the contact your st	Blood pressure  PRN MEDICA  Acetaminop  Dimenhydri  Diphenhydr  Epinephrine severe anap  Hydrocortise  Oxygen via  Salbutamol  Other (indice)  PRN medicate  regarding informa  I have rea	e meds on hold ATIONS FOR ohen 325-650 mg amine 25-50 mg (1:1000) 0.01 ohylactic reactione 100 mg IV mask/nasal pro 2 puffs q 4-6 hotate):ations not requition shared on t d, understand, escription is an origin	I: YES NO Please specify:  INFUSION REACTIONS  ING PO PRN q 4-6 hours, for pain and fever, chills INFO/IV PRN q 4 hours, for nausea and vomiting INFO/IV PRN q 4 hours, for nausea and vomiting INFO/IV PRN q 4-6 hours for itching, urticaria, prunchly (max. 0.5 mL) SC/IM PRN q 10-15 minutes × 2 con INFO PRN × 1 for severe allergic/anaphylactic reaction longs PRN for shortness of breath, wheezing ours via aerochamber PRN for dyspnea, wheezing  INFO Please specify:  INFO PRIVATE OF THE	NO e reverse.  MM/YYYY  MM/YYYY
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion Diphenhydramine 50 mg PO 15-30 min pre infusion Methylprednisolone 100 mg IV in 50 mL 0.9% sodium chloride injection, USP 15-30 min prior to infusion Other (indicate): Pre-medications not required Please specify:  PHYSICIAN SIGNATURE Do you accept that Pfizer Canada's Drug Safety Unit contact you r Notes:  SIGN HERE: † Effective date. Order(s) expires one year from the date of signature. Prescriber certificat  PATIENT SIGNATURE  SIGN HERE:  I have read and understood the Patient Consent text printed of accordance with these terms.	Blood pressure  PRN MEDICA  Acetaminop  Dimenhydri  Diphenhydri  Epinephrine severe anap  Hydrocortise  Oxygen via  Salbutamol  Other (indice  PRN medicates  regarding informates  I have readition: I certify that this present the back of this	e meds on hold ATIONS FOR ohen 325-650 mg amine 25-50 mg (1:1000) 0.01 ohylactic reaction one 100 mg IV mask/nasal pro 2 puffs q 4-6 he ate):  ations not requition shared on the d, understand, escription is an original of the form and agrees a form and agrees of the same of the sam	I: YES NO Please specify:  INFUSION REACTIONS  ING PO PRN q 4-6 hours, for pain and fever, chills IPO/IV PRN q 4 hours, for nausea and vomiting ING PO/IV/IM PRN q 4-6 hours for itching, urticaria, prunchly (max. 0.5 mL) SC/IM PRN q 10-15 minutes × 2 con  PRN × 1 for severe allergic/anaphylactic reaction longs PRN for shortness of breath, wheezing ours via aerochamber PRN for dyspnea, wheezing  Irred Please specify:  Ithis form or any accompanying document? YES Or and agree to the physician consent statement on the pate of the physician	NO e reverse.  MM/YYYY mation in
into an infusion bag containing either 0.9% Sodium Chloride Injection USP or 5% Dextrose Injection USP  PRE-MEDICATION ORDER  Acetaminophen 650 mg PO 15-30 min pre infusion  Diphenhydramine 50 mg PO 15-30 min pre infusion  Methylprednisolone 100 mg IV in 50 mL  0.9% sodium chloride injection, USP 15-30 min prior to infusion  Other (indicate):  Pre-medications not required  Please specify:  PHYSICIAN SIGNATURE  Do you accept that Pfizer Canada's Drug Safety Unit contact your Notes:  SIGN HERE:  † Effective date. Order(s) expires one year from the date of signature. Prescriber certificated PATIENT SIGNATURE  SIGN HERE:  I have read and understood the Patient Consent text printed of	Blood pressure  PRN MEDICA  Acetaminop  Dimenhydri  Diphenhydri  Epinephrine severe anap  Hydrocortise  Oxygen via  Salbutamol  Other (indice)  PRN medicate  regarding informat  I have readion: I certify that this present the back of this grinformation and	e meds on hold ATIONS FOR When 325-650 m Inate 25-50 mg It is in the control of t	I: YES NO Please specify:  INFUSION REACTIONS  ING PO PRN q 4-6 hours, for pain and fever, chills INFO/IV PRN q 4 hours, for nausea and vomiting INFO/IV PRN q 4 hours, for nausea and vomiting INFO/IV PRN q 4-6 hours for itching, urticaria, prunchly (max. 0.5 mL) SC/IM PRN q 10-15 minutes × 2 con INFO PRN × 1 for severe allergic/anaphylactic reaction longs PRN for shortness of breath, wheezing ours via aerochamber PRN for dyspnea, wheezing  INFO Please specify:  INFO PROVIDE TO PR	NO e reverse.  MM/YYYY mation in e providers











## Fax completed form to 1-833-958-3539 or 1-833-958-FLEX

#### **PATIENT CONSENT**

#### Agreement to Disclose Personal Information - PfizerFlex Program

Special Instructions: This consent form may contain words or phrases that are new to you. If any part of this form is not clear to you, please ask the person who gave you this form to explain it to you. Words that are written in **bold type** are explained on the bottom of this section.

We are asking for your permission to collect, to use and to share your Personal Information\*. The patient assistance program for RUXIENCE®, called PfizerFlex† ("Program") is a free Program offered to all patients who have been prescribed RUXIENCE®. The Program can help you in a number of ways. Sharing your Personal Information as described on this form will help us figure out which Program services and materials are best for you.

For you to take part in the Program and for us to carry out the Program activities for you, you agree to:

- Allow your **Healthcare Providers**<sup>†</sup>, the Administrators (the service providers elected by Pfizer to administer the program offerings) and the PfizerFlex Program Personnel® ("Program Personnel") to collect, use, share with each other, and store your Personal Information. These people are described at the bottom of this form.
- Allow the Program Personnel to use the Personal Information that you provide to contact you, and to collect other Personal Information from you that is needed or related to the administration of the Program. For example, this may include, asking for your feedback on the quality of the services offered by the Program or any other related services, or your progress while taking the medication RUXIENCE®, and may include limited market research, such as surveys on your experiences, so that Pfizer may better understand and improve its products and programs. Program Personnel may leave messages for you at the phone number you give them, if you have checked the can leave a message box on this enrolment form.
- Allow Pfizer Canada (the company that sells RUXIENCE®) and its affiliates ("Pfizer") to collect your Personal Information and information on any unwanted drug effects ("adverse drug events", or side effects) that you may have while taking RUXIENCE®, or other medications made by Pfizer. Commonly, Pfizer and Health Canada (the government body that approves the use of this and other medications) ask for this information to track the safety record of these medications. The information collected from you and others taking these medications allows them to better understand how these medications can affect the patients who take them. This information may be provided to Health Canada or to another regulatory agency to report any adverse drug events, or as otherwise may be required by law. Pfizer may also contact your Healthcare Providers if they need more information.
- Allow Pfizer, or a service provider hired by Pfizer, to have access to your Personal Information in order to audit the Program or provide recommendations on how to improve the Program. For example, Pfizer or its service provider may review documents that contain your Personal Information, or monitor phone conversations between you and Program Personnel for quality control purposes. Any service provider will be required to only use your Personal Information for purposes relating to the audit/Program administration, and will not disclose your Personal Information to third parties.
- The Administrator and Program Personnel can administer the prescribed RUXIENCE® medication to me during a pre-scheduled specialty clinic appointment. Such treatment shall include administration of prescribed pre-medication and management of infusion related reactions or emergencies during the infusion treatment appointment.

By giving your consent, you understand that:

- You agree to receive Program services, support and materials suitable for your needs.
- The Program Personnel are not allowed to collect, use, share or store your Personal Information

for anything other than the activities described in this consent form. They cannot share any of your Personal Information with anyone other than your Healthcare Providers, unless the Health Information\* that identifies you is removed. For example, your name, address and any personal identifiers must be removed if any of your Health Information is shared with anyone who is not your Healthcare Provider. Health Information which does not have your name, address or personal identifiers could still be shared after you withdraw your consent.

- You may take back your consent at any time by calling the Administrators at 1-855-935-FLEX (3539) or sending a request with your signature to the Administrators by fax to 1-833-958-FLEX (3539). Your consent is needed to receive services from the PfizerFlex Program. If you decide to take back your consent, you will no longer be enrolled in the PfizerFlex Program. This means that you will not be able to receive any support services from the Program, and you may not be able to get financial assistance for RUXIENCE® if you are eligible.
- · Except where prohibited by law, you may have a copy of your Personal Information. You can correct any mistakes and/or ask the Administrators any questions about the collection, use, sharing and storage of your Personal Information. You may contact the Administrators by calling 1-855-935-FLEX (3539) or by faxing your request to 1-833-958-FLEX (3539).
- Any calls to or from the Administrators while providing services of the Program may be monitored or recorded for control of quality and to train their personnel.
- Your Personal Information may be collected, used, shared and/or stored outside of your province or territory or country. The laws of those countries regarding privacy may be less strict than the laws of Canada and its provinces.
- · Your Personal Information may also be disclosed and/or transferred to a third party in the event of a proposed or actual purchase, sale (including a liquidation, realization, foreclosure or repossession), lease, amalgamation or any other type of acquisition, disposal, transfer, conveyance or financing of all or any portion of Pfizer Canada or of any of the business or assets or shares of Pfizer Canada or a division thereof.
- Pfizer Canada has the right to modify or cancel the Program and the services offered by the Program at any time without prior notice to you.
- If at any time and for any reason Pfizer Canada appoints new Program Administrators, you will allow the transfer of your Personal Information by the Administrators or by Pfizer to the new Administrators in order to continue your participation in the Program.
- You will not seek to have the amount of support you receive by way of this program counted in any Government out-of-pocket expenses for prescription drugs.
- Unless your consent is withdrawn, your consent is valid for as long as you receive services from the Program and for a reasonable time thereafter.
- \* Your Personal Information includes your individual information (name, gender, address, phone number, date of birth, etc.), your financial information and your Health Information (medical history, medical condition(s), information relating to your treatment, and information relating to your health insurance, etc.).
- The PfizerFlex Program is sponsored by Pfizer Canada to help patients get access to RUXIENCE®, and to help them manage their treatment plan for the indications approved by Health Canada.
- Healthcare Providers include all of your doctors, nurses, pharmacists or pharmacy support staff, private insurance company(s), public payer(s) and any other healthcare provider or payer that may possess the necessary information.
- § PfizerFlex Program Personnel include the employees and consultants of the Administrators, as well as any service providers that are engaged by the Administrators to manage or perform Program services and activities.

## **PHYSICIAN CONSENT**

### My signature acknowledges that:

- I am the prescribing physician of this patient;
- I have prescribed this patient RUXIENCE® for a Health Canada approved indication;
- Subject to the above-noted patient's consent and only to the extent of such patient's consent:
  - I consent to the PfizerFlex Program Personnel® contacting me with regard to the above-noted patient to assist it in administering the program, and without limitation with regard to patient reimbursement, and patient care;
  - I consent to the Administrators (the service providers elected by Pfizer to administer the program offerings) receiving, collecting, storing, using and disclosing any of my information that I provide in respect to the patient that is necessary to assist the patient in obtaining any services or assistance the patient has authorized and consented to;
  - I consent to Pfizer Canada (the company who sells RUXIENCE®) and its affiliates ("Pfizer") to contact

- me with regard to the above-noted patient if they require further information on adverse drug event pertaining to RUXIENCE®, or other medications manufactured by Pfizer;
- I agree to allow the Administrators to provide this prescription to the pharmacy chosen by the above-named patient or another pharmacy (where applicable) to ensure the patient obtains access to the therapy I have prescribed;
- I agree to allow the Administrators to contact me for any other information regarding the PfizerFlex Program\*\* that would result in enhancing the delivery or the quality of services offered by this program to my patient.
- ¶ PfizerFlex Program Personnel include the employees and consultants of the Administrators elected by Pfizer to administer the Program.
- \*\* The **PfizerFlex Program** is sponsored by Pfizer Canada to help patients get access to RUXIENCE®, and to help them manage their treatment plan for the indications approved by Health Canada.

For more information, please refer to the RUXIENCE® Product Monograph.

The Product Monograph is available upon request or it can be accessed at http://pfizer.ca/pm/en/RUXIENCE.pdf.









